

Borel Complexity of Archimedean Orders

Antoine Poulin

April 19, 2022

Left-Orderable groups

Throughout, Γ is an infinite group.

Definition

A total order on a group Γ is a **left-order** if

$$g < h \Rightarrow kg < kh.$$

We say that Γ is **left-orderable** if there is some left-order on it.

Examples

- $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$ are all left-orderable groups.
- If H, K are left-orderable groups and

$$1 \rightarrow K \rightarrow \Gamma \rightarrow H \rightarrow 1$$

Γ is left-orderable.

- $\Gamma = \langle x, y \mid yxy^{-1} = x^{-1} \rangle$ is a semi-direct product $\mathbb{Z} \rtimes \mathbb{Z}$, hence left-orderable.

Different Characterization

Definition

We say that $P \subset \Gamma$ is a **positive cone** if

- $P \cdot P \subset P$
- $P \sqcup P^{-1} = \Gamma - \{1\}$

Proposition

If $<$ is a left-order on Γ ,

$$P_{<} := \{g \in \Gamma : g > \text{id}\}$$

is a positive cone.

Duality between Orders and Cones

On the other hand, if P is a positive cone, we get a left-order :

$$g <_P h \Leftrightarrow g^{-1}h \in P.$$

Proposition

- $P = P_{<_P}$
- $g < h \Leftrightarrow g <_{P_{<}} h$

Definition of $\text{LO}(\Gamma)$

Since left-orders and positive cones are interchangeable,

Definition

The **space of left-orders** of Γ is defined as

$$\text{LO}(\Gamma) := \left\{ P \in 2^\Gamma : P \text{ is a positive cone} \right\}$$

If Γ is countable, this is compact Polish.

Definition of Archimedean Orders

Definition

We say that an order is **Archimedean** if there is for any $g, h \in P_{<}$,

$$\exists n, h < g^n$$

We extend this definition to positive cones. We also have a Polish space of Archimedean order, $\text{Ar}(\Gamma)$.

Examples

- $\mathbb{Z}, \mathbb{Q}, \mathbb{R}$ are Archimedean-orderable.
- $\Gamma = \langle x, y \mid yxy^{-1} = x^{-1} \rangle$ admits no Archimedean order.

Definition of Isomorphism Action

Definition

The **isomorphism action** $\text{Aut}(\Gamma) \curvearrowright \text{LO}(\Gamma)$ is defined by

$$\phi \cdot P := \phi(P) = \{\phi(h) : h \in P\}.$$

This restricts to an action

$$\text{Aut}(\Gamma) \curvearrowright \text{Ar}(\Gamma)$$

Motivation

We are interested in the complexity of $\text{GL}(\mathbb{Z}^2) \curvearrowright \text{Ar}(\mathbb{Z}^2)$,
motivated by

Theorem, Calderoni-Marker-Motto Ros-Shani

$\text{GL}(\mathbb{Q}^2) \curvearrowright \text{Ar}(\mathbb{Q}^2)$ is not smooth.

Still not known whether it is hyperfinite.

What does $\text{Ar}(\mathbb{Z}^2)$ look like ?

Theorem, folklore

If $P \in \text{LO}(\mathbb{Z}^2)$, there is a line Δ such that $\mathbb{R}^2 - \Delta$ has one component with only positive elements and one component with only negative elements.

If $P \in \text{Ar}(\mathbb{Z}^2)$, $\Delta \cap \mathbb{Z}^2 = \emptyset$.

The big picture




Figure – A positive cone in \mathbb{Z}^2 . Bigger dots represent elements of P and the shaded region is the half-plane containing only positive elements.

Consequences

There is a 2-1 map from $\text{Ar}(\mathbb{Z}^2)$ to line Δ which do not intersect \mathbb{Z}^2 . This is equivalent to having $\begin{pmatrix} \alpha \\ 1 \end{pmatrix} \in \Delta$, where α is irrational.

We can act as if the map is 1-1, since in each preimage we can pick canonically the cone with $\begin{pmatrix} 0 \\ 1 \end{pmatrix} \in P$.

Definition

Definition

The **action by Möbius transformations** is the action $\text{GL}(\mathbb{Z}^2) \curvearrowright \text{Irr}$ defined by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \alpha := \frac{a\alpha + b}{c\alpha + d}$$

Action on lines

$$\text{If } \begin{pmatrix} \alpha \\ 1 \end{pmatrix} \in \Delta,$$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \begin{pmatrix} \alpha \\ 1 \end{pmatrix} \in \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \Delta$$

$$\Rightarrow \begin{pmatrix} a\alpha + b \\ c\alpha + d \end{pmatrix} \in \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \Delta$$

$$\Rightarrow \begin{pmatrix} \frac{a\alpha+b}{c\alpha+d} \\ 1 \end{pmatrix} \in \begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot \Delta$$

Recap

- The action $\text{GL}(\mathbb{Z}^2) \curvearrowright \text{Ar}(\mathbb{Z}^2)$ is bireducible to the action $\text{GL}(\mathbb{Z}^2)$ on lines with irrational slope.
- The action $\text{GL}(\mathbb{Z}^2)$ on lines with irrational slope is bireducible with the action by Möbius transformations.

Continuous fractions

There is an homeomorphism $\text{Irr} \cong \mathbb{N}^{\mathbb{N}}$ defined by

$$[a_0, a_1, \dots] := a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \dots}}$$

Generators for $\text{GL}(\mathbb{Z}^2)$

We know that $\text{GL}(\mathbb{Z}^2)$ is generated by

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

These matrices act nicely through Möbius transformations.

First two matrices

We have that

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \cdot [a_0, a_1, \dots] = \begin{cases} [a_1, a_2, a_3, \dots] & \text{if } a_0 = 0 \\ [0, a_0, a_1, \dots] & \text{if } a_0 \neq 0 \end{cases}$$

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \cdot [a_0, a_1, \dots] = [a_0 + 1, a_1, \dots]$$

Chaining these two matrices, we can get tail equivalence relation

$$[a_0, \dots, a_n, c_0, c_1, \dots] \sim [b_0, \dots, b_m, c_0, c_1, \dots]$$

Last Matrix

What about $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$?

Theorem, noted in Jackson-Kechris-Louveau, proof found in Hardy-Wright

The orbit equivalence of Möbius transformations on $\mathbb{Z} \times \mathbb{N}^{\mathbb{N}}$ is exactly tail equivalence relation.

Theorem

Theorem[P.]

The isomorphism relation $\text{GL}(\mathbb{Z}^2) \curvearrowright \text{Ar}(\mathbb{Z}^2)$ is hyperfinite, but not smooth.

Question[Calderoni-Marker-Motto Ros-Shani]

How complicated is $\text{GL}(\mathbb{Q}^n) \curvearrowright \text{Ar}(\mathbb{Q}^n)$?

Higher dimensions

Theorem[P.]

The isomorphism relation $\text{GL}(\mathbb{Z}^n) \curvearrowright \text{Ar}(\mathbb{Z}^n)$ is not treeable for $n \geq 4$.

Proof based on work of Popa and Vaes.

Thank you !